



2015 INFLUENZA VACCINE WAIVER/DECLINATION

Influenza vaccine is **STRONGLY RECOMMENDED FOR HEALTHCARE WORKERS**, not only to protect themselves, but to reduce the chance of spreading influenza to our patients and community. Influenza infection can lead to serious complications and can be fatal, especially in elderly or sick persons, including those who are hospitalized. When infection occurs despite vaccination, it is usually milder.

QUESTION	Yes	No
1. Have you had a severe (life threatening) allergic reaction to any component of the vaccine including egg protein or to a previous dose of any influenza vaccination?		
2. Do you have a history of allergy to eggs? <i>If yes, please consult with your physician before receiving the vaccine.</i>		
3. Do you have a history of Guillain-Barre syndrome (a severe paralytic illness, also called GBS) that has occurred within 6 weeks of receipt of a prior influenza vaccine? <i>If yes, please consult with your physician before receiving the vaccine.</i>		

IF YOU HAVE ANSWERED YES TO ANY QUESTIONS LISTED ABOVE, PROCEED TO WAIVER OF VACCINE SECTION.

WAIVER

Complete if not eligible to receive vaccine

I am not eligible to receive the influenza vaccine today based on reason(s) marked above. I understand that I will be required to wear a surgical mask within six feet of a patient when engaged in patient care or having contact with patients while performing assigned duties for the duration of the respiratory virus season, which is generally October through March.

Signature : _____

Date: _____

DECLINATION

I am eligible to receive the influenza vaccine, BUT DO NOT WANT to take it. I understand that by refusing the vaccine I may be putting my SELF, FAMILY, and PATIENTS at risk of getting influenza. I am aware that hospitalized patients are at increased risk of getting serious complications following influenza infection.

I am declining receipt of flu vaccine based on reasons of conscience, including religious beliefs. I understand that I will be required to wear a surgical mask within six feet of a patient when engaged in patient care or having contact with patients while performing assigned duties for the duration of the respiratory virus season, which is generally October through March.

Name: _____

Signature : _____

Date: _____

